

Antibiotics  
in PID

# Use of antibiotics in the treatment of primary immunodeficiencies (PIDs)

hello@piduk.org  
0800 987 8986  
[www.piduk.org](http://www.piduk.org)

## About this booklet

Antibiotics are important medicines for managing and treating infections caused by bacteria. This booklet for patients with a primary immunodeficiency (PID) and their carers aims to promote the safe and effective use of antibiotics. It has been produced by the PID UK Medical Advisory Panel and Patient Representative Panel to help answer the questions patients and their families may have about the use of antibiotics in the treatment of PIDs but should not replace advice from a clinical immunologist.

## Contents

What are antibiotics?	3
What types of antibiotics are available?	3
Which antibiotic should you take?	3
How to take antibiotics	4
The do's and don'ts	4
What are the possible complications of taking antibiotics?	5
Interactions with other medications	5
How is the use of antibiotics in PID different?	6
About prophylactic antibiotics	7
Glossary of terms	8

## What are antibiotics?

Antibiotics are medicines used to treat infections caused by bacteria. They work either by killing the bacteria or by slowing their growth. Antibiotics are used in primary immunodeficiencies (PIDs):

- To treat bacterial infections, such as chest, sinus or ear infections
- As prophylaxis, which is prolonged treatment with the aim of preventing bacterial infection, or reducing its frequency or severity.

## What types of antibiotics are available?

There are many different types of antibiotics that are used to treat different kinds of bacterial infections. Antibiotics are often categorised by their 'spectrum' of activity.

A 'broad-spectrum' antibiotic is one that acts against a wide variety of bacteria. Examples of broad-spectrum antibiotic tablets often used in PIDs are co-amoxiclav, doxycycline and azithromycin.

Other antibiotics work against certain types of bacteria and are used only in very specific circumstances, such as when laboratory test results identify a specific bacteria.

## Which antibiotic should you take?



Your doctor will decide which antibiotic is appropriate for you based on your clinical history, examination and laboratory results (e.g. which bacteria grow from a sputum sample). In addition, each hospital will have local guidance on antibiotic use to support your doctor in the decision-making process.

Please remember to always give a sputum sample, where possible, before taking antibiotics for a chest infection. This might help your doctor make decisions on antibiotic treatment now and in the future.

## How to take antibiotics

Antibiotic treatment is usually taken by mouth in tablet form (or suspensions for children). Antibiotics can also be administered into a vein (intravenous) or for certain infections applied directly to the affected part of the body (topical; e.g. antibiotic eye drops). Occasionally and for very specific lung infections, inhaled (nebulised) antibiotics may be used.

## The do's and don'ts

If your doctor has decided that antibiotics are necessary, it is very important to take them in a responsible manner.

- Follow the instructions on the label or patient information sheet. Some antibiotic tablets are best taken on an empty stomach (an hour before or two hours after meals), whilst others should not be taken with certain foods.
- Make sure you take the antibiotics at the correct time and complete the course. Not completing the course might contribute to bacteria becoming 'antibiotic resistant' and the antibiotic being less effective in the future. This is particularly important in immunodeficiency because antibiotics can sometimes be used regularly.
- You should tell your doctor if your symptoms continue to get worse, you are unable to take the prescribed antibiotics, or you experience unacceptable side effects.

Sometimes your doctor may decide that intravenous antibiotics are needed. Intravenous antibiotics are often used for the most severe infections. However, certain antibiotics do not work very well (or at all) when given orally. In these situations intravenous antibiotics are necessary. Hospital admission for intravenous antibiotics is often required, but longer-term administration can sometimes be facilitated at home.

### Remember

You should always complete the course of antibiotics, even if you start to feel better.

## What are the possible complications of taking antibiotics?



Antibiotics are safe and highly effective medications, but as with all therapies, there are sometimes unwanted effects (side effects).

Nausea, vomiting and diarrhoea are the most common side effects. Experiencing one or more of these side effects may mean that the particular antibiotic you

have been prescribed is not suitable for you at this time. You should contact the prescribing team for further advice on what to do next.

Occasionally antibiotics lead to a fungal infection, or 'thrush' of the mouth, digestive tract or vagina. This is rarely serious and is readily treated with antifungal drops and creams where necessary.

Allergy to antibiotics is rare. Antibiotics can lead to rashes, swelling of the skin or tongue and difficulty breathing. If you have had a reaction to an antibiotic in the past, it is important to remind your doctor, nurse or pharmacist of this because your reaction may have an impact on the types of antibiotics you are given in the future.

### Important

Inform your doctor immediately if you get a reaction while on an antibiotic. If you are unable to contact your doctor, stop the antibiotic and continue to try and contact your medical team.

## Interactions with other medications

Some antibiotics can affect how other medications work. For example, some make the oral contraceptive pill less effective, in which case you may need to use a different form of contraception. Some may affect the activity of blood-thinning medication, such as warfarin, or tablets used to control epilepsy. Your doctor, nurse or pharmacist will be happy to help you with any concerns.

## How is the use of antibiotics in PID different?

The use of antibiotics in PID may differ from general use in a number of ways. Occasionally it may be necessary to remind a healthcare professional of these differences. This may be particularly important when dealing with new or temporary staff, as they may not have your full medical details to hand.

The main differences are:

### **Longer courses and stronger courses**

Patients with PID are often prescribed longer and stronger courses of antibiotics than usual. This might mean courses of 14 days, or sometimes longer. The reason for this is to prevent relapse or recurrence of infection. As always, you must complete the course, even if you are feeling better.

### **Standby antibiotics at home**

Under certain circumstances a 'reserve' course of antibiotics to be kept at home may be recommended. Your doctor will give you very specific advice regarding when these antibiotics are to be used. It can be helpful to collect a sputum sample (in the case of a chest infection) before these emergency antibiotics are started. This can be dropped into your family practice. You should keep a record of when you use these standby antibiotics, and obtain a replacement supply after they have been used.

### **Long-term (prophylactic) antibiotics** (see page 7)

Antibiotics are often prescribed for much longer periods of time with the aim of reducing the number and severity of infections. As with all antibiotics, it is particularly important to take prophylactic antibiotics exactly as prescribed and not to skip or miss doses.

### **Four golden rules of taking antibiotics in PID**

- Always take your antibiotics exactly as directed.
- Always complete the course.
- Inform your doctor, nurse or pharmacist if you experience side effects.
- Seek advice from your medical team if you do not feel better or you think things are getting worse.

## About prophylactic antibiotics

Long-term antibiotics are often used in PID to reduce the number and severity of infections. They can be used as a standalone treatment or be added to other therapies, such as immunoglobulin replacement. While much of what we know about antibiotic prophylaxis is taken from studies in conditions such as cystic fibrosis and bronchiectasis, this treatment approach has been found to be very effective by doctors specialising in PID and is common practice.

The choice and dose of prophylactic antibiotic depends on the type of PID; the presence of complications such as lung, sinus or ear disease; information from previous laboratory tests; and local guidelines on antibiotic use. Patient-specific concerns, such as dosing intervals and previous side effects, are also taken into account when making a selection.

Your doctor will carefully weigh up the risks and benefits of using antibiotic prophylaxis before offering this treatment. How well this treatment strategy works will be an important part of your regular PID follow-up. If you and your doctor find no discernible benefit from long-term antibiotics then they may be changed to an alternative or discontinued entirely. Prophylactic antibiotics will never be prescribed without good reason.

Prophylactic antibiotics are well tolerated and problems with their use are not common. Concerns have been raised about the impact of certain antibiotic families on the heart when they are used for long periods. This may be particularly relevant for older patients and those with known heart problems. If your doctor is considering the use of such medications, he or she may wish to obtain an ECG (heart trace) and discuss the risks with you before starting this approach.

Antibiotic resistance is a concern for doctors and patients alike. Studies in other diseases do suggest that long-term use of certain antibiotic drugs can contribute to a level of resistance, but this has not been well studied in PID. Some specialists will change a patient's antibiotics every few months in an attempt to avoid resistance, but the benefit of this approach is uncertain.

At present, the best advice to minimise resistance is to take the antibiotic regime exactly as prescribed by your team, never skip doses, and raise any concerns you have with your specialist directly.

## Glossary of terms

**allergy** – an exaggerated sensitivity resulting from a heightened or altered reactivity of the immune system to an external substance.

**antifungal** – a medicine used to treat infections caused by fungi.

**bronchiectasis** – a widening of the tubes (bronchi) that lead to the air sacs of the lung; this can happen because of repeated bouts of infections.

**clinical history** – the patient’s ‘story’ of their medical information.

**clinical immunologist** – a doctor that specialises in looking after patients who have problems with their immune system.

**ECG** – electrocardiogram. A way of detecting abnormal heart rhythms and investigating the cause of chest pains.

**epilepsy** – any disorder that results in seizures.

**intravenous** – inside or into a vein; e.g. antibiotics may be given directly into a vein.

**prophylactic/prophylaxis** – something that works to defend or protect against disease.

**sinuses** – air-filled space within the bones of the face and around the nose. Infection of the sinuses is called sinusitis.

**sputum** – mucus that is coughed up from your airways.

## Notes

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Notes

Lined writing area with horizontal lines.

Notes

Lined writing area with horizontal lines.

# About Primary Immunodeficiency UK

**Primary Immunodeficiency UK  
(PID UK) is a national organisation  
supporting individuals and families affected  
by primary immunodeficiencies (PIDs).**

We are the UK national member of the International Patient Organisation for Primary Immunodeficiencies (IPOPI), an association of national patient organisations dedicated to improving awareness, access to early diagnosis and optimal treatments for PID patients worldwide.

Our website at [www.piduk.org](http://www.piduk.org) provides useful information on a range of conditions and topics, and explains the work we do to ensure the voice of PID patients is heard.

If we can be of any help, please contact us at [hello@piduk.org](mailto:hello@piduk.org) or on **0800 987 8986** where you can leave a message.

Support us by becoming a member of PID UK. It's free and easy to do via our website at [www.piduk.org/register](http://www.piduk.org/register) or just get in touch with us. Members get monthly bulletins.

PID UK is reliant on voluntary donations. To make a donation, please go to [www.piduk.org/donate](http://www.piduk.org/donate)

[hello@piduk.org](mailto:hello@piduk.org)  
0800 987 8986  
[www.piduk.org](http://www.piduk.org)



This booklet was made possible, in part, by a grant provided by the Jeffrey Modell Foundation WIN Program.